NaProTechnology Success rates (in percent)

Polycystic ovarian disease

### **Artificial Reproductive Technologies** Success rates (in percent)

Polycystic ovarian disease

	TO THE OID I K	LOIMING		
Creighton Model FertilityCare System Perfect use 99.5 Typical use 96.8		Birth control pills 99.5 Perfect use 90 – 96 Typical use		
	Infertility Ti	REATMENT		
NaProTechnology Endometriosis Polycystic ovaries	$56.7 - 76.4^{3}$ $62.5 - 80.0^{3}$	21.2 <sup>3</sup> 25.6 <sup>3</sup>	n vitro fertilization Endometriosis Polycystic ovaries	
Tubal occlusion	38.43	$27.2^{3}$	Tubal occlusion	
Surgical NaProTechnology a Endometriosis	ssociated with 56.7 – 76.4 <sup>2</sup>	57.0 <sup>2</sup>	rad. surgical approach Endometriosis	(rarely used)

To Avoid Pregnancy

# DIAGNOSIS OF LUTEAL PHASE

 $41.8^{2}$ 

NaProTechnology Current medical approach Detect by properly targeting  $98.6^{7}$ Not available n/a hormone evaluation

 $62.5 - 80.0^{2}$ 

### Premenstrual Dysphoric Disorder (PMS)

 $95.2^{4}$ NaProTechnology Current treatment 43.0 Antidepressants

#### Postpartum Depression

Slow improvement over 6–12 months NaProTechnology  $92.4 - 96.7^{5}$ Generally within 1-30 days anti-anxiety meds

#### PREMATURITY & SEVERE PREMATURITY RATE

NaProTechnology		T	raditional treatment
Prematurity rate	$7.0^{6}$	12.9	Prematurity rate
Severe prematurity rate	$1.3^{6}$	3.9	Severe prematurity rate

## RECURRENT SPONTANEOUS ABORTION

NaProTechnology 79.0 Lower Current medical approach

## Dating the Beginning of Pregnancy

 $100.0^{8}$ NaProTechnology 86.08 Using date of last menstrual period

#### CHRONIC PELVIC PAIN

Surgical NaProTechnology	decreased 2.4x	Current medical approach
Hysterectomy rate	11.5	40.0 Hysterectomy rate


### Cost-effectiveness

Creighton Model System	\$494 <sup>9</sup>	\$1,866°	Birth control pills
Infertility	\$32210	$$9,226^{10}$	IVF
Prematurity	\$16,79511	\$28,55611	Current medical approach
PMS evaluation & treatment	\$3,21812	$$5,104^{12}$	Current medical approach

- 1. Completely comparable to oral contraceptives.
- 2. Measured by survival curve analysis at 36 months, compared to published results from Johns Hopkins University Medical Center.
- 3. A range of effectiveness acquired from different study designs.
- and oral naltrexone.
- 5. With the use of IM progesterone therapy.
- 6. Using the Prematurity Prevention Protocol of the Pope Paul VI Institute.
- 7. Using the Creighton Model FertilityCare System to target evaluation of the post-ovulatory hormone phase of the cycle.
- Within 10 days.
- 9. Based on 5 years of use.
- 10. Based on costs per cycle of treatment.
- 4. With the use of targeted HCG hormonal support 11. Based on cost saving generated by decrease in prematurity rate to 7.0 percent
  - 12. Includes cost savings due to improved productivity.

This table produced by Pope Paul VI Institute for the Study of Human Reproduction, Omaha, NE 2004. NaProTechnology is a registered trademark of the Pope Paul VI Institute for the Study of Human Reproduction. It can be freely used by any person or entity so long as its use reflects the medical concepts and values expressed in the textbook The Medical & Surgical Practice of NaProTechnology.





For the Study of Human Reproduction

6901 Mercy Rd. Omaha, NE 68106 (402) 390-6600

www.popepaulvi.com